

Random square-tiled surfaces and random multicurves in large genus

Anton Zorich

(after joint works with V. Delecroix, E. Goujard and P. Zograf)

Surfaces, Manifolds and Related Topics
Luxembourg, March 20, 2024

Count of square-tiled surfaces

- Square-tiled surfaces: formal definition

- Intersection numbers

- Volume polynomials

- Ribbon graphs

- Count of metric ribbon graphs

- Multicurve associated to a square-tiled surface

- Ribbon graph decomposition of a square-tiled surface

- Count of square-tiled surfaces: an algorithm

- Surface decompositions

- Associated polynomials

- Volume of \mathcal{Q}_2

Mirzakhani's count of closed geodesics

Random square-tiled surfaces

Count of square-tiled surfaces. (Masur–Veech volume of the moduli space of quadratic differentials)

A square-tiled surface



Square-tiled surfaces: formal definition

Take a finite set of copies of identical oriented squares for which two opposite sides are chosen to be horizontal and the remaining two sides are declared to be vertical. Identify pairs of sides of the squares by isometries in such way that horizontal sides are glued to horizontal sides and vertical sides to vertical. We get a topological surface S without boundary. We consider only those surfaces obtained in this way which are connected and oriented. The form dz^2 on each square is compatible with the gluing and endows S with a complex structure and with a non-zero quadratic differential $q = dz^2$ with at most simple poles. We call such a surface a *square-tiled surface*.

Square-tiled surfaces: formal definition

Take a finite set of copies of identical oriented squares for which two opposite sides are chosen to be horizontal and the remaining two sides are declared to be vertical. Identify pairs of sides of the squares by isometries in such way that horizontal sides are glued to horizontal sides and vertical sides to vertical. We get a topological surface S without boundary. We consider only those surfaces obtained in this way which are connected and oriented. The form dz^2 on each square is compatible with the gluing and endows S with a complex structure and with a non-zero quadratic differential $q = dz^2$ with at most simple poles. We call such a surface a *square-tiled surface*.

Fix the genus g of the surface and the number n of corners with cone angle π (the ones adjacent to exactly two squares). The question on the asymptotic number of such square-tiled surfaces tiled with at most $N \gg 1$ squares (connected covers over \mathbb{CP}^1 ramified over 4 points and having prescribed ramification profile) is equivalent to evaluation of the Masur–Veech volume of the moduli space of quadratic differentials.

Intersection numbers (Witten–Kontsevich correlators)

The Deligne–Mumford compactification $\overline{\mathcal{M}}_{g,n}$ of the moduli space of smooth complex curves of genus g with n labeled marked points $P_1, \dots, P_n \in C$ is a complex orbifold of complex dimension $3g - 3 + n$.

Choose index i in $\{1, \dots, n\}$. The family of complex lines cotangent to C at the point P_i forms a holomorphic line bundle \mathcal{L}_i over $\mathcal{M}_{g,n}$ which extends to $\overline{\mathcal{M}}_{g,n}$. The first Chern class of this *tautological bundle* is denoted by $\psi_i = c_1(\mathcal{L}_i)$.

Any collection of nonnegative integers satisfying $d_1 + \dots + d_n = 3g - 3 + n$ determines a positive rational “*intersection number*” (or the “*correlator*” in the physical context):

$$\langle \tau_{d_1} \dots \tau_{d_n} \rangle_g := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n} .$$

Intersection numbers (Witten–Kontsevich correlators)

The Deligne–Mumford compactification $\overline{\mathcal{M}}_{g,n}$ of the moduli space of smooth complex curves of genus g with n labeled marked points $P_1, \dots, P_n \in C$ is a complex orbifold of complex dimension $3g - 3 + n$.

Choose index i in $\{1, \dots, n\}$. The family of complex lines cotangent to C at the point P_i forms a holomorphic line bundle \mathcal{L}_i over $\mathcal{M}_{g,n}$ which extends to $\overline{\mathcal{M}}_{g,n}$. The first Chern class of this *tautological bundle* is denoted by $\psi_i = c_1(\mathcal{L}_i)$.

Any collection of nonnegative integers satisfying $d_1 + \dots + d_n = 3g - 3 + n$ determines a positive rational “*intersection number*” (or the “*correlator*” in the physical context):

$$\langle \tau_{d_1} \dots \tau_{d_n} \rangle_g := \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n} .$$

The famous Witten’s conjecture claims that these numbers satisfy certain recurrence relations which are equivalent to certain differential equations on the associated generating function (“*partition function in 2-dimensional quantum gravity*”). Witten’s conjecture was proved by M. Kontsevich; alternative proofs belong to A. Okounkov and R. Pandharipande, to M. Mirzakhani, to M. Kazarian and S. Lando (and there are more).

Volume polynomials

Consider the moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n marked points. Let d_1, \dots, d_n be an ordered partition of $3g - 3 + n$ into the sum of nonnegative numbers, $d_1 + \dots + d_n = 3g - 3 + n$, let \mathbf{d} be the multiindex (d_1, \dots, d_n) and let $b^{2\mathbf{d}}$ denote $b_1^{2d_1} \dots b_n^{2d_n}$.

Define the homogeneous polynomial $N_{g,n}(b_1, \dots, b_n)$ of degree $6g - 6 + 2n$ in variables b_1, \dots, b_n :

$$N_{g,n}(b_1, \dots, b_n) := \sum_{|\mathbf{d}|=3g-3+n} c_{\mathbf{d}} b^{2\mathbf{d}},$$

where

$$c_{\mathbf{d}} := \frac{1}{2^{5g-6+2n} \mathbf{d}!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}$$

Volume polynomials

Consider the moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n marked points. Let d_1, \dots, d_n be an ordered partition of $3g - 3 + n$ into the sum of nonnegative numbers, $d_1 + \dots + d_n = 3g - 3 + n$, let \mathbf{d} be the multiindex (d_1, \dots, d_n) and let $b^{2\mathbf{d}}$ denote $b_1^{2d_1} \dots b_n^{2d_n}$.

Define the homogeneous polynomial $N_{g,n}(b_1, \dots, b_n)$ of degree $6g - 6 + 2n$ in variables b_1, \dots, b_n :

$$N_{g,n}(b_1, \dots, b_n) := \sum_{|\mathbf{d}|=3g-3+n} c_{\mathbf{d}} b^{2\mathbf{d}},$$

where

$$c_{\mathbf{d}} := \frac{1}{2^{5g-6+2n} \mathbf{d}!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}$$

Up to a numerical factor, the polynomial $N_{g,n}(b_1, \dots, b_n)$ coincides with the top homogeneous part of the Mirzakhani's volume polynomial $V_{g,n}(b_1, \dots, b_n)$ providing the Weil–Petersson volume of the moduli space of bordered Riemann surfaces:

$$V_{g,n}^{top}(b) = 2^{2g-3+n} \cdot N_{g,n}(b).$$

Volume polynomials

Consider the moduli space $\mathcal{M}_{g,n}$ of Riemann surfaces of genus g with n marked points. Let d_1, \dots, d_n be an ordered partition of $3g - 3 + n$ into the sum of nonnegative numbers, $d_1 + \dots + d_n = 3g - 3 + n$, let \mathbf{d} be the multiindex (d_1, \dots, d_n) and let $b^{2\mathbf{d}}$ denote $b_1^{2d_1} \dots b_n^{2d_n}$.

Define the homogeneous polynomial $N_{g,n}(b_1, \dots, b_n)$ of degree $6g - 6 + 2n$ in variables b_1, \dots, b_n :

$$N_{g,n}(b_1, \dots, b_n) := \sum_{|\mathbf{d}|=3g-3+n} c_{\mathbf{d}} b^{2\mathbf{d}},$$

where


$$c_{\mathbf{d}} := \frac{1}{2^{5g-6+2n} \mathbf{d}!} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \dots \psi_n^{d_n}$$

Define the formal operation \mathcal{Z} on monomials as

$$\mathcal{Z} : \prod_{i=1}^n b_i^{m_i} \longmapsto \prod_{i=1}^n (m_i! \cdot \zeta(m_i + 1)),$$

and extend it to symmetric polynomials in b_i by linearity.

Trivalent ribbon graphs



This trivalent ribbon graph defines an orientable surface of genus $g = 1$ with $n = 2$ boundary components. Assigning lengths to all edges of the core graph, we endow each boundary component with an induced length defined as the sum of the lengths of edges which it follows.

Note, however, that in general, fixing a genus g , a number n of boundary components and integer lengths b_1, \dots, b_n of boundary components, we get plenty of trivalent integral metric ribbon graphs associated to such data. The Theorems of Kontsevich and Norbury count them.

Count of metric ribbon graphs

Theorem (Kontsevich'92; in this stronger form — Norbury'10). Consider a collection of positive integers b_1, \dots, b_n such that $\sum_{i=1}^n b_i$ is even. The weighted count of genus g connected trivalent metric ribbon graphs Γ with integer edges and with n labeled boundary components of lengths b_1, \dots, b_n is equal to $N_{g,n}(b_1, \dots, b_n)$ up to the lower order terms:

$$\sum_{\Gamma \in \mathcal{R}_{g,n}} \frac{1}{|\text{Aut}(\Gamma)|} N_{\Gamma}(b_1, \dots, b_n) = N_{g,n}(b_1, \dots, b_n) + \text{lower order terms},$$

where $\mathcal{R}_{g,n}$ denote the set of (nonisomorphic) trivalent ribbon graphs Γ of genus g and with n boundary components.

Count of metric ribbon graphs

Theorem (Kontsevich'92; in this stronger form — Norbury'10). Consider a collection of positive integers b_1, \dots, b_n such that $\sum_{i=1}^n b_i$ is even. The weighted count of genus g connected trivalent metric ribbon graphs Γ with integer edges and with n labeled boundary components of lengths b_1, \dots, b_n is equal to $N_{g,n}(b_1, \dots, b_n)$ up to the lower order terms:


$$\sum_{\Gamma \in \mathcal{R}_{g,n}} \frac{1}{|\text{Aut}(\Gamma)|} N_{\Gamma}(b_1, \dots, b_n) = N_{g,n}(b_1, \dots, b_n) + \text{lower order terms},$$

where $\mathcal{R}_{g,n}$ denote the set of (nonisomorphic) trivalent ribbon graphs Γ of genus g and with n boundary components.

(Formal statement justifying the notion of “lower order terms”: the right-hand side is a *quasipolynomial* in the integers b_1, \dots, b_n depending on the number k of odd b_i . The top homogeneous part is zero when k is odd.)

A version of this Theorem is an important part of Kontsevich's proof of Witten's conjecture.


Multicurve associated to a square-tiled surface



Having a square-tiled surface we associate to it a topological surface S on which we mark all “corners” with cone angle π (i.e. vertices with exactly two adjacent squares). By convention the associated hyperbolic metric has cusps at the marked points.

We also consider a multicurve γ on the resulting surface composed of the waist curves γ_j of all maximal horizontal cylinders. We encode the number of horizontal bands of squares in each cylinder by taking the components of the multicurve with integer weights.

Ribbon graph decomposition of a square-tiled surface



Leaves of the horizontal foliation on the square-tiled surface passing through singular points (in blue) are called *critical*. Considering tubular neighborhoods of these critical leaves we get metric ribbon graphs. Cylinders (represented by the multicurve in red) are joining boundary components of these ribbon graphs.

A dual graph to the multicurve is called *stable graph* Γ . The vertices of Γ are in the natural bijection with metric ribbon graphs given by components of $S \setminus \gamma$. The edges are in the bijection with the waist curves γ_i of the cylinders. The marked points are encoded by “legs” — half-edges of the dual graph.


Count of square-tiled surfaces: an algorithm

1. Fix a genus g and a number n of corners (conical points) of angle π .
2. Consider a finite collection of *stable graphs* encoding all possible admissible decompositions of a hyperbolic surface of genus g with n cusps (equivalently, all complex stable curves of genus g with n marked points).
3. For each stable graph with k edges associate formal variables b_1, \dots, b_k to its edges and associate metric ribbon graphs to the vertices.
4. Using the Kontsevich–Norbury count of metric ribbon graphs, count the number of ways to join them by square-tiled cylinders.


Count of square-tiled surfaces: an algorithm

1. Fix a genus g and a number n of corners (conical points) of angle π .
2. Consider a finite collection of *stable graphs* encoding all possible admissible decompositions of a hyperbolic surface of genus g with n cusps (equivalently, all complex stable curves of genus g with n marked points).
3. For each stable graph with k edges associate formal variables b_1, \dots, b_k to its edges and associate metric ribbon graphs to the vertices.
4. Using the Kontsevich–Norbury count of metric ribbon graphs, count the number of ways to join them by square-tiled cylinders.



Masur–Veech volume of the moduli space of quadratic differentials. This moduli space is the total space of the cotangent bundle to the moduli space $\mathcal{M}_{g,n}$ of complex curves with n marked points and, hence, has a canonical symplectic structure, and a volume element. Square-tiled surfaces represent integer points in this space. Those, which are tiled with at most N squares, are integer points in a “bundle of balls of radius N ” over $\mathcal{M}_{g,n}$. Thus, asymptotics of the number of square-tiled surfaces of genus g with n conical points of angle π tiled with at most $N \rightarrow +\infty$ squares gives us the Masur–Veech volume.





$$\frac{1}{2} \cdot 1 \cdot b_1 \cdot N_{1,2}(b_1, b_1)$$





$$\frac{1}{2} \cdot \frac{1}{2} \cdot b_1 \cdot N_{1,1}(b_1) \cdot N_{1,1}(b_1)$$





$$\frac{1}{8} \cdot 1 \cdot b_1 b_2 \cdot N_{0,4}(b_1, b_1, b_2, b_2)$$




$$\frac{1}{2} \cdot \frac{1}{2} \cdot b_1 b_2 \cdot N_{0,3}(b_1, b_1, b_2) \cdot N_{1,1}(b_2)$$




$$\frac{1}{8} \cdot \frac{1}{2} \cdot b_1 b_2 b_3 \cdot N_{0,3}(b_1, b_1, b_2) \cdot N_{0,3}(b_2, b_3, b_3)$$




$$\frac{1}{12} \cdot \frac{1}{2} \cdot b_1 b_2 b_3 \cdot N_{0,3}(b_1, b_2, b_3) \cdot N_{0,3}(b_1, b_2, b_3)$$







$$b_1 \quad \frac{1}{2} \cdot 1 \cdot b_1 \cdot N_{1,2}(b_1, b_1) = \frac{1}{2} \cdot b_1 \left(\frac{1}{384} (2b_1^2)(2b_1^2) \right)$$




$$b_1 \quad \frac{1}{2} \cdot \frac{1}{2} \cdot b_1 \cdot N_{1,1}(b_1) \cdot N_{1,1}(b_1) = \frac{1}{4} \cdot b_1 \left(\frac{1}{48} b_1^2 \right) \left(\frac{1}{48} b_1^2 \right)$$




$$b_1 \quad b_2 \quad \frac{1}{8} \cdot 1 \cdot b_1 b_2 \cdot N_{0,4}(b_1, b_1, b_2, b_2) = \frac{1}{8} \cdot b_1 b_2 \cdot \left(\frac{1}{4} (2b_1^2 + 2b_2^2) \right)$$



$$b_1 \quad b_2 \quad \frac{1}{2} \cdot \frac{1}{2} \cdot b_1 b_2 \cdot N_{0,3}(b_1, b_1, b_2) \cdot N_{1,1}(b_2) = \frac{1}{4} \cdot b_1 b_2 \cdot (1) \cdot \left(\frac{1}{48} b_2^2 \right)$$




$$b_1 \quad b_2 \quad b_3 \quad \frac{1}{8} \cdot \frac{1}{2} \cdot b_1 b_2 b_3 \cdot N_{0,3}(b_1, b_1, b_2) \cdot N_{0,3}(b_2, b_3, b_3) = \frac{1}{16} \cdot b_1 b_2 b_3 \cdot (1) \cdot (1)$$




$$b_1 \quad b_2 \quad b_3 \quad \frac{1}{12} \cdot \frac{1}{2} \cdot b_1 b_2 b_3 \cdot N_{0,3}(b_1, b_2, b_3) \cdot N_{0,3}(b_1, b_2, b_3) = \frac{1}{24} \cdot b_1 b_2 b_3 \cdot (1) \cdot (1)$$


Volume of \mathcal{Q}_2




$$b_1 \quad \frac{1}{192} \cdot b_1^5 \xrightarrow{\mathcal{Z}} \frac{1}{192} \cdot (5! \cdot \zeta(6)) = \frac{1}{1512} \cdot \pi^6$$




$$b_1 \quad \frac{1}{9216} \cdot b_1^5 \xrightarrow{\mathcal{Z}} \frac{1}{9216} \cdot (5! \cdot \zeta(6)) = \frac{1}{72576} \cdot \pi^6$$



$$b_1 \quad b_2 \quad \frac{1}{16} (b_1^3 b_2 + b_1 b_2^3) \xrightarrow{\mathcal{Z}} \frac{1}{16} \cdot 2(1! \cdot \zeta(2)) \cdot (3! \cdot \zeta(4)) = \frac{1}{720} \cdot \pi^6$$



$$b_1 \quad b_2 \quad \frac{1}{192} \cdot b_1 b_2^3 \xrightarrow{\mathcal{Z}} \frac{1}{192} \cdot (1! \cdot \zeta(2)) \cdot (3! \cdot \zeta(4)) = \frac{1}{17280} \cdot \pi^6$$



$$b_1 \quad b_2 \quad b_3 \quad \frac{1}{16} b_1 b_2 b_3 \xrightarrow{\mathcal{Z}} \frac{1}{16} \cdot (1! \cdot \zeta(2))^3 = \frac{1}{3456} \cdot \pi^6$$





$$b_1 \quad b_2 \quad b_3 \quad \frac{1}{24} b_1 b_2 b_3 \xrightarrow{\mathcal{Z}} \frac{1}{24} \cdot (1! \cdot \zeta(2))^3 = \frac{1}{5184} \cdot \pi^6$$


$$\text{Vol } \mathcal{Q}_2 = \frac{128}{5} \cdot \left(\frac{1}{1512} + \frac{1}{72576} + \frac{1}{720} + \frac{1}{17280} + \frac{1}{3456} + \frac{1}{5184} \right) \cdot \pi^6 = \frac{1}{15} \pi^6.$$


Volume of \mathcal{Q}_2


$$b_1 \text{  \quad \frac{1}{192} \cdot b_1^5 \xrightarrow{\mathcal{Z}} \frac{1}{192} \cdot (5! \cdot \zeta(6)) = \frac{1}{1512} \cdot \pi^6$$

$$\text{ \quad \frac{1}{9216} \cdot b_1^5 \xrightarrow{\mathcal{Z}} \frac{1}{9216} \cdot (5! \cdot \zeta(6)) = \frac{1}{72576} \cdot \pi^6$$

$$b_1 \text{  b_2 \quad \frac{1}{16} (b_1^3 b_2 + b_1 b_2^3) \xrightarrow{\mathcal{Z}} \frac{1}{16} \cdot 2(1! \cdot \zeta(2)) \cdot (3! \cdot \zeta(4)) = \frac{1}{720} \cdot \pi^6$$

$$b_1 \text{  b_2 \quad \frac{1}{192} \cdot b_1 b_2^3 \xrightarrow{\mathcal{Z}} \frac{1}{192} \cdot (1! \cdot \zeta(2)) \cdot (3! \cdot \zeta(4)) = \frac{1}{17280} \cdot \pi^6$$

$$b_1 \text{  b_3 \quad \frac{1}{16} b_1 b_2 b_3 \xrightarrow{\mathcal{Z}} \frac{1}{16} \cdot (1! \cdot \zeta(2))^3 = \frac{1}{3456} \cdot \pi^6$$

$$b_1 \text{  b_3 \quad \frac{1}{24} b_1 b_2 b_3 \xrightarrow{\mathcal{Z}} \frac{1}{24} \cdot (1! \cdot \zeta(2))^3 = \frac{1}{5184} \cdot \pi^6$$

$$\text{Vol } \mathcal{Q}_2 = \frac{128}{5} \cdot \left(\frac{1}{1512} + \frac{1}{72576} + \frac{1}{720} + \frac{1}{17280} + \frac{1}{3456} + \frac{1}{5184} \right) \cdot \pi^6 = \frac{1}{15} \pi^6.$$

Count of square-tiled
surfaces

**Mirzakhani's count of
closed geodesics**

- Multicurves
- Geodesic
representatives of
multicurves
- Frequencies of
multicurves
- Example
- Hyperbolic and flat
geodesic multicurves


Random square-tiled
surfaces

Mirzakhani's count of simple closed geodesics

Simple closed multicurve, its topological type and underlying primitive multicurve


Having an arbitrary collection of complicated non self-intersecting and non pairwise intersecting curves (called a *multicurve*), one can apply an appropriate diffeomorphism of the surface which “unwraps” the multicurve to a simple canonical representative.

A general multicurve ρ :




the canonical representative $\gamma = 3\gamma_1 + \gamma_2 + 2\gamma_3$ in its orbit $\text{Mod}_2 \cdot \rho$ under the action of the mapping class group and the associated *reduced* multicurve.

$$\gamma = 3\gamma_1 + \gamma_2 + 2\gamma_3.$$




$$\gamma_{\text{reduced}} = \gamma_1 + \gamma_2 + \gamma_3$$



(You can practice in unwinding curves at <https://aharalab.sakura.ne.jp/teruaki.html>)

Geodesic representatives of multicurves

Consider several pairwise nonintersecting essential simple closed curves $\gamma_1, \dots, \gamma_k$ on a smooth surface $S_{g,n}$ of genus g with n punctures. In the presence of a hyperbolic metric X on $S_{g,n}$ the simple closed curves $\gamma_1, \dots, \gamma_k$ contract to simple closed geodesics.



Fact. For any hyperbolic metric X the simple closed geodesics representing $\gamma_1, \dots, \gamma_k$ do not have pairwise intersections.

We define the hyperbolic length of a multicurve $\gamma := \sum_{i=1}^k a_i \gamma_i$ as $\ell_\gamma(X) := \sum_{i=1}^k a_i \ell_X(\gamma_i)$, where $\ell_X(\gamma_i)$ is the hyperbolic length of the simple closed geodesic in the free homotopy class of γ_i .

Denote by $s_X(L, \gamma)$ the number of simple closed geodesic multicurves on X of topological type $[\gamma]$ and of hyperbolic length at most L .

Frequencies of multicurves

Theorem (Mirzakhani'08). *For any integral multi-curve γ and any hyperbolic surface X in $\mathcal{M}_{g,n}$ the number $s_X(L, \gamma)$ of simple closed geodesic multicurves on X of topological type $[\gamma]$ and of hyperbolic length at most L has the following asymptotics:*

$$s_X(L, \gamma) \sim \mu_{\text{Th}}(B_X) \cdot \frac{c(\gamma)}{b_{g,n}} \cdot L^{6g-6+2n} \quad \text{as } L \rightarrow +\infty.$$

Here $\mu_{\text{Th}}(B_X)$ depends only on the hyperbolic metric X ; the constant $b_{g,n}$ depends only on g and n ; $c(\gamma)$ depends only on the topological type of γ and admits a closed formula (in terms of the intersection numbers of ψ -classes).

Frequencies of multicurves

Theorem (Mirzakhani'08). *For any integral multi-curve γ and any hyperbolic surface X in $\mathcal{M}_{g,n}$ the number $s_X(L, \gamma)$ of simple closed geodesic multicurves on X of topological type $[\gamma]$ and of hyperbolic length at most L has the following asymptotics:*

$$s_X(L, \gamma) \sim \mu_{\text{Th}}(B_X) \cdot \frac{c(\gamma)}{b_{g,n}} \cdot L^{6g-6+2n} \quad \text{as } L \rightarrow +\infty.$$



Here $\mu_{\text{Th}}(B_X)$ depends only on the hyperbolic metric X ; the constant $b_{g,n}$ depends only on g and n ; $c(\gamma)$ depends only on the topological type of γ and admits a closed formula (in terms of the intersection numbers of ψ -classes).

Corollary (Mirzakhani'08). *For any hyperbolic surface X in $\mathcal{M}_{g,n}$, and any two rational multicurves γ_1, γ_2 on a smooth surface $S_{g,n}$ considered up to the action of the mapping class group one obtains*

$$\lim_{L \rightarrow +\infty} \frac{s_X(L, \gamma_1)}{s_X(L, \gamma_2)} = \frac{c(\gamma_1)}{c(\gamma_2)}.$$


Example

A simple closed geodesic on a hyperbolic sphere with six cusps separates the sphere into two components. We either get three cusps on each of these components (as on the left picture) or two cusps on one component and four cusps on the complementary component (as on the right picture). Hyperbolic geometry excludes other partitions.



Example

A simple closed geodesic on a hyperbolic sphere with six cusps separates the sphere into two components. We either get three cusps on each of these components (as on the left picture) or two cusps on one component and four cusps on the complementary component (as on the right picture). Hyperbolic geometry excludes other partitions.




Example (Mirzakhani'08); confirmed experimentally in 2017 by M. Bell; confirmed in 2017 by more implicit computer experiment of V. Delecroix and by relating it to Masur–Veech volume.

$$\lim_{L \rightarrow +\infty} \frac{\text{Number of } (3 + 3)\text{-simple closed geodesics of length at most } L}{\text{Number of } (2 + 4)\text{-simple closed geodesics of length at most } L} = \frac{4}{3}.$$

Example


A simple closed geodesic on a hyperbolic sphere with six cusps separates the sphere into two components. We either get three cusps on each of these components (as on the left picture) or two cusps on one component and four cusps on the complementary component (as on the right picture). Hyperbolic geometry excludes other partitions.



In this sense one can say that for any hyperbolic metric X on a sphere with 6 cusps, a long simple closed geodesic separates the cusps as $(3 + 3)$ with probability $\frac{4}{7}$ and as $(2 + 4)$ with probability $\frac{3}{7}$.


$$\lim_{L \rightarrow +\infty} \frac{\text{Number of } (3 + 3)\text{-simple closed geodesics of length at most } L}{\text{Number of } (2 + 4)\text{-simple closed geodesics of length at most } L} = \frac{4}{3}.$$

Hyperbolic and flat geodesic multicurves



Left picture represents a geodesic multicurve $\gamma = 2\gamma_1 + \gamma_2 + \gamma_3 + 2\gamma_4$ on a hyperbolic surface in $\mathcal{M}_{0,7}$. Right picture represents the same multicurve this time realized as the union of the waist curves of horizontal cylinders of a square-tiled surface of the same genus, where cusps of the hyperbolic surface are in the one-to-one correspondence with the conical points having cone angle π (i.e. with the simple poles of the corresponding quadratic differential). The weights of individual connected components γ_i are recorded by the heights of the cylinders. Clearly, there are plenty of square-tiled surface realizing this multicurve.

Hyperbolic and flat geodesic multicurves




Theorem (Delecroix–Goujard–Zograf–Zorich’21). For any topological class γ of simple closed multicurves considered up to homeomorphisms of a surface $S_{g,n}$, the associated Mirzakhani’s asymptotic frequency $c(\gamma)$ of **hyperbolic** multicurves coincides with the asymptotic frequency of simple closed **flat** geodesic multicurves of type γ represented by associated square-tiled surfaces.


Remark. Francisco Arana Herrera recently found an alternative proof of this result. His proof uses more geometric approach.

Multicurves on a surface of genus two and their frequencies


The picture below illustrates all topological types of primitive multicurves on a surface of genus two without punctures; the fractions give frequencies of non-primitive multicurves γ having a reduced multicurve $\gamma_{reduced}$ of the corresponding type.




$$\frac{16}{63}$$




$$\frac{8}{15}$$




$$\frac{1}{9}$$



$$\frac{1}{189}$$



$$\frac{1}{45}$$



$$\frac{2}{27}$$

In genus 3 there are already 41 types of multicurves, in genus 4 there are 378 types, in genus 5 there are 4554 types and this number grows faster than exponentially when genus g grows. It becomes pointless to produce tables: we need to extract a reasonable sub-collection of most common types which ideally, carry all Thurston's measure when $g \rightarrow +\infty$.

Count of square-tiled
surfaces

Mirzakhani's count of
closed geodesics

Random square-tiled
surfaces

- Random integers
- Random permutations
- Shape of a random multicurve
- Heights of cylinders of a random square-tiled surface
- Main Theorem (informally)
- Keystone underlying result
- Combinatorial formulation of Witten's conjecture
- Arnold's problem
- Rue des Petits-Carreux

Shape of a random multicurve on a surface of large genus. Shape of a random square-tiled surface of large genus.

Statistics of prime decompositions: random integer numbers

The Prime Number Theorem states that an integer number n taken randomly in a large interval $[1, N]$ is prime with asymptotic probability $\frac{\log N}{N}$.

Actually, one can tell much more about prime decomposition of a large random integer. Denote by $\omega(n)$ the number of prime divisors of an integer n counted without multiplicities. In other words, if n has prime decomposition $n = p_1^{m_1} \dots p_k^{m_k}$, let $\omega(n) = k$. By the Erdős–Kac theorem, the centered and rescaled distribution prescribed by the counting function $\omega(n)$ tends to the normal distribution:

Erdős–Kac Theorem (1939)

$$\lim_{N \rightarrow +\infty} \frac{1}{N} \text{card} \left\{ n \leq N \mid \frac{\omega(n) - \log \log N}{\sqrt{\log \log N}} \leq x \right\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt.$$

The subsequent results of A. Selberg (1954) and of A. Rényi and P. Turán (1958) describe the rate of convergence.

Statistics of prime decompositions: random permutations

Denote by $K_n(\sigma)$ the number of disjoint cycles in the cycle decomposition of a permutation σ in the symmetric group S_n . Consider the uniform probability measure on S_n . A random permutation σ of n elements has exactly k cycles in its cyclic decomposition with probability $\mathbb{P}(K_n(\sigma) = k) = \frac{s(n,k)}{n!}$, where $s(n, k)$ is the unsigned Stirling number of the first kind. It is immediate to see that $\mathbb{P}(K_n(\sigma) = 1) = \frac{1}{n}$. V. L. Goncharov computed the expected value and the variance of K_n as $n \rightarrow +\infty$:



$$\mathbb{E}(K_n) = \log n + \gamma + o(1), \quad \mathbb{V}(K_n) = \log n + \gamma - \zeta(2) + o(1),$$

and proved the following central limit theorem:

Theorem (V. L. Goncharov, 1944)

$$\lim_{n \rightarrow +\infty} \frac{1}{n!} \text{card} \left\{ \sigma \in S_n \mid \frac{K_n(\sigma) - \log n}{\sqrt{\log n}} \leq x \right\} = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^x e^{-\frac{t^2}{2}} dt.$$

Shape of a random square-tiled surface of large genus




Questions.

- How many singular horizontal leaves (in blue on the right picture) has a random square-tiled surface of genus g ?
- Find the probability distribution for the number $K_g(S) = 1, 2, 3, \dots, 3g - 3$ of maximal horizontal cylinders (represented by red waist curves on the left picture)
- What are the typical heights h_1, \dots, h_k of the cylinders?
- What is the shape of a random square-tiled surface of large genus?

Shape of a random multicurve (random square-tiled surface) on a surface of large genus in simple words

Theorem (Delecroix–Goujard–Zograf–Zorich'20.) *With probability which tends to 1 as $g \rightarrow \infty$,*

- *The reduced multicurve $\gamma_{reduced} = \gamma_1 + \dots + \gamma_k$ associated to a random integral multicurve $\gamma = m_1\gamma_1 + \dots + m_k\gamma_k$ does not separate the surface;*
- *$\gamma_{reduced}$ has about $(\log g)/2$ components and has one of the following types:*




$0.09 \log(g)$ components

...

...

...

...



$0.62 \log(g)$ components

$$\mathbb{P}\left(0.09 \log g < K_g(\gamma) < 0.62 \log g\right) = 1 - O\left((\log g)^{24} g^{-1/4}\right).$$

*A random square-tiled surface (without conical points of angle π) of large genus has about $\frac{\log(g)}{2}$ cylinders, and **all conical points sit at the same horizontal and the same vertical level with probability which tends to 1 as $g \rightarrow \infty$.***

Heights of cylinders of a random square-tiled surface

Theorem (Delecroix–Goujard–Zograf–Zorich’19). *If we fix any k and consider only k -cylinder square-tiled surfaces, then a (conditional) probability that every horizontal cylinder is composed of a single band of squares tends to 1 as $g \rightarrow +\infty$.*

Theorem (Delecroix–Goujard–Zograf–Zorich’19). *If we do not fix the number of horizontal cylinders, then the probability that every horizontal cylinder of a random square-tiled surface is composed of a single band of squares tends to $\frac{\sqrt{2}}{2}$ as genus grows. More generally, each of the heights m_1, \dots, m_k of horizontal cylinders of a random square-tiled surface is bounded from above by an integer m with probability which tends to $\sqrt{\frac{m}{m+1}}$ as $g \rightarrow +\infty$.*

However, the mean value of $m_1 + \dots + m_k$ is infinite in any genus g .

Main Theorem (informally)

Main Theorem (Delecroix–Goujard–Zograf–Zorich’20). *As g grows, the probability distribution $\mathbb{P}(K_g = k)$ rapidly becomes, basically, indistinguishable from the distribution of the number $K_{3g-3}(\sigma)$ of disjoint cycles in a random permutation σ of $3g - 3$ elements (with respect to some explicit nonuniform probability measure on the symmetric group). In particular, for any $j \in \mathbb{N}$ the difference of the j -th moments of the two distributions is of the order $O(g^{-1})$. We have an explicit asymptotic formula for all cumulants. It gives*

$$\mathbb{E}(K_g) = \frac{\log(6g - 6)}{2} + \frac{\gamma}{2} + \log 2 + o(1),$$
$$\mathbb{V}(K_g) = \frac{\log(6g - 6)}{2} + \frac{\gamma}{2} + \log 2 - \frac{3}{4}\zeta(2) + o(1),$$

where $\gamma = 0.5772\dots$ denotes the Euler–Mascheroni constant.

In practice, already for $g = 12$ the match of the graphs of the distributions is such that they are visually indistinguishable.

Keystone underlying result

Our results use the Delecroix–Goujard–Zograf–Zorich’19 conjecture proved in

Theorem (Aggarwal’21). *The Masur–Veech volume of the moduli space of holomorphic quadratic differentials has the following large genus asymptotics:*

$$\text{Vol } \mathcal{Q}_g \sim \frac{4}{\pi} \cdot \left(\frac{8}{3}\right)^{4g-4} \quad \text{as } g \rightarrow +\infty.$$

Keystone underlying result

Our results use the Delecroix–Goujard–Zograf–Zorich’19 conjecture proved in **Theorem (Aggarwal’21)**. *The Masur–Veech volume of the moduli space of holomorphic quadratic differentials has the following large genus asymptotics:*

$$\text{Vol } \mathcal{Q}_g \sim \frac{4}{\pi} \cdot \left(\frac{8}{3}\right)^{4g-4} \quad \text{as } g \rightarrow +\infty.$$

The similar conjecture of Eskin–Zorich’03 on the large genus asymptotics of Masur–Veech volumes of individual strata of *Abelian* differentials is recently proved by Aggarwal’19 and by Chen–Möller–Sauvaget–Zagier’20. The analogous conjecture for *quadratic* differentials still resists:

Conjecture (ADGZZ’20). *The Masur–Veech volume of any stratum of meromorphic quadratic differentials with at most simple poles has the following large genus asymptotics (with the error term uniformly small for all partitions \mathbf{d}):*

$$\text{Vol } \mathcal{Q}(d_1, \dots, d_n) \stackrel{?}{\sim} \frac{4}{\pi} \cdot \prod_{i=1}^n \frac{2^{d_i+2}}{d_i + 2} \quad \text{as } g \rightarrow +\infty,$$

under assumption that the number of simple poles is bounded or grows much slower than the genus.

Combinatorial formulation of Witten's conjecture

Initial data: $\langle \tau_0^3 \rangle = 1, \quad \langle \tau_1 \rangle = \frac{1}{24}.$

String equation:

$$\langle \tau_0 \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n+1} = \langle \tau_{d_1-1} \cdots \tau_{d_n} \rangle_{g,n} + \cdots + \langle \tau_{d_1} \cdots \tau_{d_n-1} \rangle_{g,n}.$$

Dilaton equation:

$$\langle \tau_1 \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n+1} = (2g - 2 + n) \langle \tau_{d_1} \cdots \tau_{d_n} \rangle_{g,n}.$$

Virasoro constraints (in Dijkgraaf–Verlinde–Verlinde form; $k \geq 1$):

$$\begin{aligned} \langle \tau_{k+1} \tau_{d_1} \cdots \tau_{d_n} \rangle_g &= \frac{1}{(2k+3)!!} \left[\sum_{j=1}^n \frac{(2k+2d_j+1)!!}{(2d_j-1)!!} \langle \tau_{d_1} \cdots \tau_{d_j+k} \cdots \tau_{d_n} \rangle_g \right. \\ &\quad + \frac{1}{2} \sum_{\substack{r+s=k-1 \\ r,s \geq 0}} (2r+1)!! (2s+1)!! \langle \tau_r \tau_s \tau_{d_1} \cdots \tau_{d_n} \rangle_{g-1} \\ &\quad \left. + \frac{1}{2} \sum_{\substack{r+s=k-1 \\ r,s \geq 0}} (2r+1)!! (2s+1)!! \sum_{\{1,\dots,n\}=I \amalg J} \langle \tau_r \prod_{i \in I} \tau_{d_i} \rangle_{g'} \langle \tau_s \prod_{i \in J} \tau_{d_i} \rangle_{g-g'} \right]. \end{aligned}$$

Keystone underlying result

We also strongly use the uniform large genus asymptotics of ψ -classes, which we conjectured in 2019. We proved it for 2-correlators; a general formula was proved by A. Aggarwal:


Theorem (Aggarwal'21). *The following **uniform** asymptotic formula is valid:*

$$\begin{aligned} \int_{\overline{\mathcal{M}}_{g,n}} \psi_1^{d_1} \cdots \psi_n^{d_n} &= \\ &= \frac{1}{24^g} \cdot \frac{(6g - 5 + 2n)!}{g! (3g - 3 + n)!} \cdot \frac{d_1! \cdots d_n!}{(2d_1 + 1)! \cdots (2d_n + 1)!} \cdot (1 + \varepsilon(\mathbf{d})), \end{aligned}$$

where $\varepsilon(\mathbf{d}) = O\left(1 + \frac{(n + \log g)^2}{g}\right)$ **uniformly** for all $n = o(\sqrt{g})$ and all partitions \mathbf{d} , $d_1 + \cdots + d_n = 3g - 3 + n$, as $g \rightarrow +\infty$.

Arnold's problem (2002-8)

Glue randomly two boundary components of a braid with a large number N of strands on a surface of genus $g - 1$ so that the endpoints fit.



Theorem. *The probability p_g to get a single connected curve upon a random gluing of a random braid is*

$$p_g = \frac{1}{(4g - 2)2^{2g-4} \text{Vol } \mathcal{H}_g} \rightarrow \frac{1}{4g} + o\left(\frac{1}{g}\right) \text{ as } g \rightarrow +\infty.$$

Examples: $p_1 = \frac{6}{\pi^2}$, $p_2 = \frac{45}{2\pi^4}$, $p_3 = \frac{243}{2\pi^6}$.

2^e ARR^t

RUE DES
PETITS CARRÉAUX