

Drinfeld cusp forms and their combinatorics

Gebhard Böckle
gebhard.boeckle@uni-due.de

Department of Mathematics
Universität Duisburg-Essen
45117 Essen, Germany

Workshop on
Computations with Modular Forms
Bristol, August 22, 2008

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

- Definition
- Basic properties
- The residue map

How to understand the quotient tree?

References

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Outline

G. Böckle

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

- Definition
- Basic properties
- The residue map

How to understand the quotient tree?

References

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

Outline

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

Outline

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Outline

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Basic notation

\mathbb{F}_q the field of $q = p^n$ elements

K_∞ a local field with residue field \mathbb{F}_q

\mathcal{O}_∞ the ring of integers of K_∞

π a uniformizer of K_∞

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

The Bruhat-Tits tree

G. Böckle

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q$$

Definition (Bruhat-Tits tree)

\mathcal{T} := the simplicial complex of dimension 1 with

set of vertices $\text{Vert}(\mathcal{T}) :=$

homothety classes $[L]$ of rank 2 \mathcal{O}_∞ -lattices $L \subset K_\infty^2$

set of edges $\text{Edge}(\mathcal{T}) :=$

pairs $([L], [L'])$ such that $\pi L \subsetneq L' \subsetneq L$.

$|\mathcal{T}|$ the geometric realization of \mathcal{T}

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

The Bruhat-Tits tree

G. Böckle

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q$$

Definition (Bruhat-Tits tree)

\mathcal{T} := the simplicial complex of dimension 1 with

set of vertices $\text{Vert}(\mathcal{T}) :=$

homothety classes $[L]$ of rank 2 \mathcal{O}_∞ -lattices $L \subset K_\infty^2$

set of edges $\text{Edge}(\mathcal{T}) :=$

pairs $([L], [L'])$ such that $\pi L \subsetneq L' \subsetneq L$.

$|\mathcal{T}|$ the geometric realization of \mathcal{T}

Lemma

\mathcal{T} is a $q+1$ -regular tree.

Definition

Vertices $\Lambda_i := [\mathcal{O}_\infty \oplus \pi^i \mathcal{O}_\infty]$, $i \in \mathbb{Z}$.

standard vertex Λ_0 , standard edge $e_0 := (\Lambda_0, \Lambda_1)$

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Group action

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q$$

Consider elements of K_∞^2 as **column vectors**
 \Rightarrow have natural left action of $GL_2(K_\infty)$ on K_∞^2 .

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q$$

Consider elements of K_∞^2 as **column vectors**
 \Rightarrow have natural left action of $GL_2(K_\infty)$ on K_∞^2 .

Definition ($GL_2(K_\infty)$ -action on \mathcal{T})

$$GL_2(K_\infty) \times \mathcal{T} \rightarrow \mathcal{T} : (\gamma, [L]) \mapsto [\gamma L]$$

$$\text{Set } \Gamma_\infty := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in GL_2(\mathcal{O}_\infty) \mid c \in \pi \mathcal{O}_\infty \right\}.$$

Lemma

$GL_2(K_\infty)$ acts transitively on $\text{Vert}(\mathcal{T})$ and $\text{Edge}(\mathcal{T})$.

$$\text{Vert}(\mathcal{T}) = GL_2(K_\infty) / GL_2(\mathcal{O}_\infty) K_\infty^*,$$

$$\text{Edge}(\mathcal{T}) = GL_2(K_\infty) / \Gamma_\infty K_\infty^*.$$

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Drinfeld's upper half plane Ω

$$\mathbb{C}_\infty := \widehat{K_\infty^{\text{alg}}}$$

Definition (Drinfeld's upper half plane)

$$\Omega := \mathbb{P}^1(\mathbb{C}_\infty) \setminus \mathbb{P}^1(K_\infty)$$

Definition ($GL_2(K_\infty)$ -action on Ω)

$$GL_2(K_\infty) \times \Omega \rightarrow \Omega : (\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}, z) \mapsto \gamma z = \frac{az+b}{cz+d}$$

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

(rigid) analysis on Ω

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q, \mathbb{C}_\infty, \Omega$$

Let $b_1, \dots, b_q \in \mathcal{O}_\infty$ be representatives of \mathcal{O}_∞/π .

Proposition (reduction map)

\exists a (natural) $GL_2(K_\infty)$ -equivariant map

$$\rho: \Omega \rightarrow |\mathcal{T}| \text{ such that}$$

$$\rho^{-1}(|e_0| \setminus \{\Lambda_0, \Lambda_1\}) = \{z \in \mathbb{C}_\infty \mid 1 < |z| < q\}$$

$$\rho^{-1}(\Lambda_0) = \{z \in \mathbb{C}_\infty \mid \forall_{i=1, \dots, q} |z - b_i| = 1\}$$

(rigid) analysis on Ω

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q, \mathbb{C}_\infty, \Omega$$

Let $b_1, \dots, b_q \in \mathcal{O}_\infty$ be representatives of \mathcal{O}_∞/π .

Proposition (reduction map)

\exists a (natural) $GL_2(K_\infty)$ -equivariant map

$$\rho: \Omega \rightarrow |\mathcal{T}| \text{ such that}$$

$$\rho^{-1}(|e_0| \setminus \{\Lambda_0, \Lambda_1\}) = \{z \in \mathbb{C}_\infty \mid 1 < |z| < q\}$$

$$\rho^{-1}(\Lambda_0) = \{z \in \mathbb{C}_\infty \mid \forall_{i=1, \dots, q} |z - b_i| = 1\}$$

Remarks “ Ω is like a tubular neighborhood of \mathcal{T} ”

(rigid) analysis on Ω

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q, \mathbb{C}_\infty, \Omega$$

Let $b_1, \dots, b_q \in \mathcal{O}_\infty$ be representatives of \mathcal{O}_∞/π .

Proposition (reduction map)

\exists a (natural) $GL_2(K_\infty)$ -equivariant map

$$\rho: \Omega \rightarrow |\mathcal{T}| \text{ such that}$$

$$\rho^{-1}(|e_0| \setminus \{\Lambda_0, \Lambda_1\}) = \{z \in \mathbb{C}_\infty \mid 1 < |z| < q\}$$

$$\rho^{-1}(\Lambda_0) = \{z \in \mathbb{C}_\infty \mid \forall_{i=1, \dots, q} |z - b_i| = 1\}$$

Remarks “ Ω is like a tubular neighborhood of \mathcal{T} ”
 $GL_2(K_\infty)$ -translates of $\rho^{-1}(|e_0|)$ provide an atlas for Ω .

(rigid) analysis on Ω

G. Böckle

$$K_\infty, \mathcal{O}_\infty, \pi, \mathbb{F}_q, \mathbb{C}_\infty, \Omega$$

Let $b_1, \dots, b_q \in \mathcal{O}_\infty$ be representatives of \mathcal{O}_∞/π .

Proposition (reduction map)

\exists a (natural) $GL_2(K_\infty)$ -equivariant map

$$\rho: \Omega \rightarrow |\mathcal{T}| \text{ such that}$$

$$\rho^{-1}(|e_0| \setminus \{\Lambda_0, \Lambda_1\}) = \{z \in \mathbb{C}_\infty \mid 1 < |z| < q\}$$

$$\rho^{-1}(\Lambda_0) = \{z \in \mathbb{C}_\infty \mid \forall_{i=1, \dots, q} |z - b_i| = 1\}$$

Remarks “ Ω is like a tubular neighborhood of \mathcal{T} ”

$GL_2(K_\infty)$ -translates of $\rho^{-1}(|e_0|)$ provide an atlas for Ω .

On these charts use Laurent series type expansions to define

(rigid) analytic functions on Ω .

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Drinfeld modular forms

From now on: $K_\infty := \mathbb{F}_q((\frac{1}{T}))$, $\pi := \frac{1}{T}$.

For $A := \mathbb{F}_q[T]$ and $K = \text{Frac}(A)$ have

$$GL_2(A) \hookrightarrow GL_2(K) \hookrightarrow GL_2(K_\infty).$$

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition
Basic properties
The residue map

How to understand the quotient tree?

References

Drinfeld modular forms

G. Böckle

From now on: $K_\infty := \mathbb{F}_q((\frac{1}{T}))$, $\pi := \frac{1}{T}$.

For $A := \mathbb{F}_q[T]$ and $K = \text{Frac}(A)$ have

$$GL_2(A) \hookrightarrow GL_2(K) \hookrightarrow GL_2(K_\infty).$$

For $\Gamma \subset GL_2(A)$ a congruence subgroup:

Definition (Drinfeld modular form (Goss))

A Drinfeld modular form of weight k (and trivial type) for Γ is a rigid analytic function

$$f: \Omega \rightarrow \mathbb{C}_\infty$$

such that

(a) $f\left(\begin{pmatrix} a & b \\ c & d \end{pmatrix} z\right) = (cz + d)^k f(z)$ for all $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$.

(b) f has a Laurent series expansion at all cusps with vanishing principal part.

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition
Basic properties
The residue map

How to understand the quotient tree?

References

Results

One defines cusp forms (in the obvious way).

Have Hecke operators for prime ideals $0 \neq \mathfrak{p} \subset \mathbb{F}_q[T]$

No known analog of a Petersson inner product.

Let f be a Hecke eigenform with eigenvalues $a_{\mathfrak{p}}(f)$.

Theorem (Goss)

The $a_{\mathfrak{p}}(f)$ are integral

$K_f := K(\{a_{\mathfrak{p}}(f)\}_{\mathfrak{p}})$ is finite over K .

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Results

One defines cusp forms (in the obvious way).

Have Hecke operators for prime ideals $0 \neq \mathfrak{p} \subset \mathbb{F}_q[T]$

No known analog of a Petersson inner product.

Let f be a Hecke eigenform with eigenvalues $a_{\mathfrak{p}}(f)$.

Theorem (Goss)

The $a_{\mathfrak{p}}(f)$ are integral

$K_f := K(\{a_{\mathfrak{p}}(f)\}_{\mathfrak{p}})$ is finite over K .

Theorem (B.)

There is a strictly compatible system

$$\left(\rho_{f,\lambda}: \text{Gal}(\overline{K}/K) \rightarrow \text{GL}_1(\widehat{K}_f^{\lambda}) \right)_{\lambda \text{ finite}}$$

such that $\rho_{f,\lambda}(\text{Frob}_{\mathfrak{p}}) = a_{\mathfrak{p}}(f)$ for almost all \mathfrak{p} .

The sequence $(a_{\mathfrak{p}}(f))_{\mathfrak{p}}$ is given by a Hecke character.

Questions

There is no multiplicity one result!

Does multiplicity one hold for fixed weight?

Does it hold in weight 2

Does it hold in weight 2 and for $\Gamma_0(\mathfrak{p})$ with \mathfrak{p} prime?

\rightsquigarrow Possible implications for uniform boundedness of torsion points of Drinfeld modules of rank 2 over K .

G. Böckle

[Drinfeld modular forms](#)

[The Bruhat-Tits tree](#)

[The Drinfeld upper half plane](#)

[Drinfeld modular forms](#)

[Harmonic cocycles](#)

[Definition](#)

[Basic properties](#)

[The residue map](#)

[How to understand the quotient tree?](#)

[References](#)

Questions

There is no multiplicity one result!

Does multiplicity one hold for fixed weight?

Does it hold in weight 2

Does it hold in weight 2 and for $\Gamma_0(p)$ with p prime?

\rightsquigarrow Possible implications for uniform boundedness of torsion points of Drinfeld modules of rank 2 over K .

There is no Ramanujan-Petersson conjecture

But each eigenvalue systems seems to have fixed weight.

What is the distribution of weights?

G. Böckle

[Drinfeld modular forms](#)

[The Bruhat-Tits tree](#)

[The Drinfeld upper half plane](#)

[Drinfeld modular forms](#)

[Harmonic cocycles](#)

[Definition](#)

[Basic properties](#)

[The residue map](#)

[How to understand the quotient tree?](#)

[References](#)

Questions

There is no multiplicity one result!

Does multiplicity one hold for fixed weight?

Does it hold in weight 2

Does it hold in weight 2 and for $\Gamma_0(p)$ with p prime?

\rightsquigarrow Possible implications for uniform boundedness of torsion points of Drinfeld modules of rank 2 over K .

There is no Ramanujan-Petersson conjecture

But each eigenvalue systems seems to have fixed weight.

What is the distribution of weights?

There may be p not dividing the level N of f such that

$$\rho_{f,\lambda}(\text{Frob}_p) \neq a_p(f)$$

(because of non-ordinariness of modular curves of level Nq)

What happens at these p ?

G. Böckle

[Drinfeld modular forms](#)

[The Bruhat-Tits tree](#)

[The Drinfeld upper half plane](#)

[Drinfeld modular forms](#)

[Harmonic cocycles](#)

[Definition](#)

[Basic properties](#)

[The residue map](#)

[How to understand the quotient tree?](#)

[References](#)

Harmonic cocycles

$\mathcal{T}, \text{Edge}(\mathcal{T}), \Gamma$

How to compute Drinfeld modular forms?

Let M be a $K[GL_2(A)]$ -module with $\dim_K(M)$ finite.

G. Böckle

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

Definition

- Basic properties
- The residue map

How to understand the quotient tree?

References

$\mathcal{T}, \text{Edge}(\mathcal{T}), \Gamma$

How to compute Drinfeld modular forms?

Let M be a $K[GL_2(A)]$ -module with $\dim_K(M)$ finite.

Definition

The K -vector space $C_{har}(\Gamma, M)$ of M -valued Γ -invariant **harmonic cocycles** is the set of maps

$$c: \text{Edge}(\mathcal{T}) \rightarrow M : e \mapsto c(e),$$

such that:

1. For all edges e one has $c(-e) = -c(e)$.
2. For all vertices v one has $\sum_{e \rightarrow v} c(e) = 0$,
where the sum is over all edges e ending at v .
3. For all $\gamma \in \Gamma$ and $e \in \text{Edge}(\mathcal{T})$ one has $c(\gamma e) = \gamma c(e)$.

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane
Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties
The residue map

How to understand the quotient tree?

References

Basic properties

\mathcal{T} , $\text{Edge}(\mathcal{T})$, Γ

Proposition (automatic cuspidality; Teitelbaum)

Given M there exists a finite subset Z of $\Gamma \backslash \mathcal{T}$ such that any $c \in C_{\text{har}}(\Gamma, M)$ vanishes on all edges e not in a class of Z .

G. Böckle

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane
Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Basic properties

G. Böckle

$\mathcal{T}, \text{Edge}(\mathcal{T}), \Gamma$

Proposition (automatic cuspidality; Teitelbaum)

Given M there exists a finite subset Z of $\Gamma \backslash \mathcal{T}$ such that any $c \in C_{\text{har}}(\Gamma, M)$ vanishes on all edges e not in a class of Z .

Definition

A simplex $t \in \text{Vert}(\mathcal{T}) \cup \text{Edge}(\mathcal{T})$ is Γ -**stable** iff

$$\text{Stab}_{\Gamma}(t) = \{1\}.$$

Proposition

There are only finitely many Γ -stable orbits of simplices.

Theorem (Teitelbaum)

Suppose Γ is p' -torsion free. Then:

- ▶ *Any Γ -invariant harmonic cocycle is determined by its values on the Γ -stable orbits of edges.*
- ▶ *The only relations are those coming from Γ -stable vertices.*

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane
Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

Remark:

The space $C_{har}(\Gamma, M)$ has an interpretation in terms of relative group homology. Let $\Gamma_v \subset \Gamma$ be the stabilizers of a set of representatives for the cusps. Then:

$$C_{har}(\Gamma, M) \cong H_1(\Gamma, \Gamma_v, M).$$

The residue map

$$\mathcal{T}, \text{Edge}(\mathcal{T}), \Gamma, C_{\text{har}}(\Gamma, M)$$

Recall:

A Drinfeld modular form f is a rigid analytic function on Ω .

Ω is a tubular neighborhood of \mathcal{T} via ρ .

G. Böckle

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

- Definition
- Basic properties
- The residue map**

How to understand the quotient tree?

References

The residue map

G. Böckle

$$\mathcal{T}, \text{Edge}(\mathcal{T}), \Gamma, C_{\text{har}}(\Gamma, M)$$

Recall:

A Drinfeld modular form f is a rigid analytic function on Ω .

Ω is a tubular neighborhood of \mathcal{T} via ρ .

ρ^{-1} of the inner part of an edge e is an annulus $A(e)$.

For f of weight 2 define

$$\text{Res}_2 : \text{Edge}(\mathcal{T}) \rightarrow \mathbb{C}_\infty : e \mapsto \text{Res}_{A(e)}(fdz).$$

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

- Definition
- Basic properties
- The residue map**

How to understand the quotient tree?

References

The residue map

G. Böckle

$$\mathcal{T}, \text{Edge}(\mathcal{T}), \Gamma, C_{\text{har}}(\Gamma, M)$$

Recall:

A Drinfeld modular form f is a rigid analytic function on Ω .

Ω is a tubular neighborhood of \mathcal{T} via ρ .

ρ^{-1} of the inner part of an edge e is an annulus $A(e)$.

For f of weight 2 define

$$\text{Res}_2: \text{Edge}(\mathcal{T}) \rightarrow \mathbb{C}_\infty : e \mapsto \text{Res}_{A(e)}(fdz).$$

Theorem (Teitelbaum)

Res₂ defines an isomorphism from the \mathbb{C}_∞ vector space of Drinfeld cusp forms of weight 2 and level Γ to

$$C_{\text{har}}(\Gamma, K) \otimes_K \mathbb{C}_\infty.$$

An analogous theorem holds in weight k with $M \approx \text{Sym}^{k-2}$.

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

- Definition
- Basic properties
- The residue map

How to understand the quotient tree?

References

On the proof of Teitelbaum's theorem:

It suffices to prove it for $\Gamma = \Gamma(N)$ with $N \in \mathbb{F}_q[T] \setminus \mathbb{F}_q$.

Injectivity: Using a π -adic measure theory, Teitelbaum constructs an explicit section for

$$\text{Res}_2: S_2(\Gamma(N)) \rightarrow C_{\text{har}}(\Gamma, \mathbb{C}_\infty).$$

Drinfeld modular forms

The Bruhat-Tits tree

The Drinfeld upper half plane

Drinfeld modular forms

Harmonic cocycles

Definition

Basic properties

The residue map

How to understand the quotient tree?

References

On the proof of Teitelbaum's theorem:

It suffices to prove it for $\Gamma = \Gamma(N)$ with $N \in \mathbb{F}_q[T] \setminus \mathbb{F}_q$.

Injectivity: Using a π -adic measure theory, Teitelbaum constructs an explicit section for

$$\text{Res}_2: S_2(\Gamma(N)) \rightarrow C_{\text{har}}(\Gamma, \mathbb{C}_\infty).$$

Surjectivity: Compute $\dim S_2(\Gamma(N))$ via Riemann-Roch and a canonical line bundle on $\Gamma(N) \backslash \Omega$.

Express $\dim C_{\text{har}}(\Gamma(N), \mathbb{C}_\infty)$ as the number of stable orbits of edges minus the number of stable orbits of vertices..

Show that the dimensions are equal.

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane
Drinfeld modular forms

Harmonic cocycles

Definition
Basic properties
The residue map

How to understand the quotient tree?

References

How to understand the quotient tree?

G. Böckle

Proposition

The quotient tree $GL_2(\mathbb{F}_q[T]) \backslash \mathcal{T}$ is represented by the half line with vertices $\{\Lambda_i\}_{i \geq 0}$.

There are no $GL_2(\mathbb{F}_q[T])$ -stable simplices of \mathcal{T} . For $i \geq 1$, the stabilizer of Λ_{i+1} is strictly larger than that of Λ_i .

Drinfeld modular forms

- The Bruhat-Tits tree
- The Drinfeld upper half plane
- Drinfeld modular forms

Harmonic cocycles

- Definition
- Basic properties
- The residue map

How to understand the quotient tree?

References

How to understand the quotient tree?

G. Böckle

Proposition

The quotient tree $GL_2(\mathbb{F}_q[T]) \backslash \mathcal{T}$ is represented by the half line with vertices $\{\Lambda_i\}_{i \geq 0}$.

There are no $GL_2(\mathbb{F}_q[T])$ -stable simplices of \mathcal{T} . For $i \geq 1$, the stabilizer of Λ_{i+1} is strictly larger than that of Λ_i .

For general Γ :

Consider $\Gamma \backslash \mathcal{T}$ as a finite ‘covering’ of the above half line.

The stabilizers of simplices of the ‘cover’ have a similar monotonicity property as those of $GL_2(\mathbb{F}_q[T]) \backslash \mathcal{T}$.

Stable simplices can only be found above Λ_i for small i (depending on Γ).

Using the above idea, one can show all ‘basic properties’ on harmonic cocycles we quoted.

Drinfeld modular forms

The Bruhat-Tits tree
The Drinfeld upper half plane
Drinfeld modular forms

Harmonic cocycles

Definition
Basic properties
The residue map

How to understand the quotient tree?

References

References

E.-U. Gekeler and U. Nonnengardt, *Fundamental domains of some arithmetic groups over function fields*, Internat. J. Math. **6** (1995), 689–708.

J.-P. Serre, *Trees*

J. Teitelbaum, *The Poisson Kernel for Drinfeld Modular Curves*, JAMS **4** (1991), No. 3, pp. 491-511.

G. Böckle

[Drinfeld modular forms](#)

- [The Bruhat-Tits tree](#)
- [The Drinfeld upper half plane](#)
- [Drinfeld modular forms](#)

[Harmonic cocycles](#)

- [Definition](#)
- [Basic properties](#)
- [The residue map](#)

[How to understand the quotient tree?](#)

[References](#)